

ECNACLYN



The study volunteers were asked to answer several questions on their satisfaction with Ecnaclyn with respect to appearance and health of the skin.

- **76% confirmed the improvement of appearance of their complexion**
- **83% were interested in further use of Ecnaclyn.**


Results of postmarketing surveillance of Ecnaclyn made by Walmark

Patients with diagnosed acne over 10 years of age were included in the study. The patients used 1 tablet of Ecnaclyn daily for 8 weeks and underwent three examinations – when entering the study, three weeks later and after eight weeks of application. At each visit the number of comedones was evaluated (closed non-inflammatory white pimples or open black comedones) and of papulopustules (inflammatory lesions – pimples). On closing the study, the patients evaluated their satisfaction with the product using a 1-5 degree scale (1 – very satisfied, 5 – unsatisfied) and also potential adverse reactions were recorded during the application. The study included 25 patients. As early as 3 weeks later, the number of both comedones and papulopustules statistically significantly decreased from the average number of 17.3 to 11.9 and from 21.6 to 9.3 respectively ($p < 0.05$) which after 8-week application further statistically significantly decreased to 5.4 and 5.2 respectively ($p < 0.05$). Final subjective evaluation of the patients' satisfaction with Ecnaclyn using a 1-5 degree scale was as follows: degree 1: 80%, degree 2: 12% and degree 3: 8%. During the product application, no study participants encountered any adverse reactions.


Conclusion

During Ecnaclyn application, the number of comedones and papulopustules in the acne patients statistically significantly decreased as early as after 3 weeks and after 8-week use it further statistically significantly dropped. According to their evaluations, 80% of the patients were very satisfied with the product.

GRAPH1: AVERAGE NUMBER OF COMEDONES AND PAPULOPUSTULES BY THE INDIVIDUAL VISITS



GRAPH 2: FINAL SUBJECTIVE EVALUATION OF THE PATIENTS' SATISFACTION WITH ECNACLYN USING A 1-5 DEGREE SCALE



PICTURE: EXAMPLES OF STUTUS BEFORE AND AFTER 8-WEEK CONSUMPTION OF ECNACLYN

A. Status before consumption

B. Status after consumption



